


Administering Vaccines: Dose, Route, Site, and Needle Size

Vaccine	Dose	Route
Diphtheria, Tetanus, Pertussis (DTaP, DT, Tdap, Td)	0.5 mL	IM
<i>Haemophilus influenzae</i> type b (Hib)	0.5 mL	IM
Hepatitis A (HepA)	≤18 yrs; 0.5 mL	IM
	≥19 yrs; 1.0 mL	
Hepatitis B (HepB) <i>*Persons 11–15 yrs may be given Recombivax HB (Merck) 1.0 mL adult formulation on a 2-dose schedule.</i>	<19yrs: 0.5 mL	IM
	≥20 yrs: 1.0 mL	
Human papillomavirus (HPV)	0.5 mL	IM
Influenza, live attenuated (LAIV)	0.2 mL	Intranasal spray
Influenza, trivalent inactivated (TIV)	6-35 mos: 0.25 mL	IM
	≥3 yrs: 0.5 mL	
TIV: Fluzone intradermal (18–64 yrs)	0.1 mL	ID
Measles, Mumps, Rubella (MMR)	0.5 mL	SC
Meningococcal – conjugate (MCV)	0.5 mL	IM
Meningococcal – polysaccharide (MPSV)	0.5 mL	SC
Pneumococcal conjugate (PCV)	0.5 mL	IM
Pneumococcal polysaccharide (PPSV)	0.5 mL	IM or SC
Polio, inactivated (IPV)	0.5 mL	IM or SC
Rotavirus (RV)	Rotarix: 1.0 mL	Oral
	Rotateq: 2.0 mL	
Varicella (Var)	0.5 mL	SC
Zoster (Zos)	0.65 mL	SC
Combination Vaccines		
DTaP-HepB-IPV (Pediarix) DTaP-IPV/Hib (Pentacel) DTaP-IPV (Kinrix) Hib-HepB (Comvax)	0.5 mL	IM
MMRV (ProQuad)	≤12 yrs: 0.5 mL	SC
HepA-HepB (Twinrix)	≥18 yrs: 1.0 mL	IM

Injection Site and Needle Size		
Subcutaneous (SC) injection Use a 23–25 gauge needle. Choose the injection site that is appropriate to the person's age and body mass.		
Age	Needle Length	Injection Site
Infants (1–12 mos)	5/8"	Fatty tissue over anterolateral thigh muscle
Children 12 mos or older, adolescents, and adults	5/8"	Fatty tissue over anterolateral thigh muscle or fatty tissue over triceps
Intramuscular (IM) injection Use a 22–25 gauge needle. Choose the injection site and needle length appropriate to the person's age and body mass.		
Age	Needle Length	Injection Site
Newborns (1 st 28 days)	5/8"*	Anterolateral thigh muscle
Infants (1–12 mos)	1"	Anterolateral thigh muscle
Toddlers (1–2 yrs)	1–1 1/4" 5/8–1"*	Anterolateral thigh muscle or deltoid muscle of arm
Children & teens (3–18 years)	5/8–1"* 1"–1 1/4"	Deltoid muscle of arm or anterolateral thigh muscle
Adults 19 yrs or older		
Male or female less than 130 lbs	5/8–1"*	Deltoid muscle of arm
Female 130–200 lbs Male 130–260 lbs	1–1 1/2"	Deltoid muscle of arm
Female 200+ lbs Male 260+ lbs	1 1/2"	Deltoid muscle of arm

*A 5/8" needle may be used for patients weighing less than 130 lbs (<60 kg) for IM injection in the deltoid muscle only if the skin is stretched tight, the subcutaneous tissue is not bunched, and the injection is made at a 90-degree angle.



Please note: Always refer to the package insert included with each biologic for complete vaccine administration information. CDC's Advisory Committee on Immunization Practices (ACIP) recommendations for the particular vaccine should be reviewed as well (see www.immunize.org/acip).