

Pediatric Vaccines

Hepatitis A & B



Disease	Vaccine	Age Group	Volume	Doses	Month Intervals
Hep A	Havrix® GlaxoSmithKline	1 - 18 years	0.5 mL	2	0, 6 - 12 months
	Vaqta® Merck	1 - 18 years	0.5 mL	2	0, 6 - 18
Hep B	Engerix-B® GlaxoSmithKline	0 - 19 years	0.5 mL	3	Birth: 0, 1 - 4, 18 Older: 0, 1 - 2, 4
	Recombivax HB® Merck	0 - 19 years	0.5 mL	3	Birth: 0, 1 - 4, 6 - 18 Older: 0, 1 - 2, 4
		11 - 15 years	1.0 mL	2	0, 4 - 6
Hib Hep B	Comvax® Merck	6 weeks - 4 years	0.5 mL	3	2, 4, 12 - 15
DTaP HepB IPV	Pediarix® GlaxoSmithKline	6 weeks - 6 years	0.5 mL	3	2, 4, 6

A birth dose of Engerix-B® or Recombivax HB® should be administered prior to discharge from a birthing hospital.

Combination Comvax® and Pediarix® vaccines cannot be administered as a birth dose, but can be used to complete the Hepatitis B vaccine series for infants, including those with HBsAg+ mothers.